

years as more information from gene sequencing databases becomes available. Functionally, these molecules are intracellular enzymes that play key roles in cell growth, differentiation and inter-cell communication. Aberrant protein kinase activity has been implicated in many disease states including several forms of cancer and severe-combined immunodeficiency disease (Barker *et al.*, 1997; Lehtola *et al.*; Arpaia *et al.* 1994; Elder *et al.* 1995; Roifman, 1995). Similarly, activation of protein kinase activity within mononuclear cells is required to drive the cytokine production which underlies many autoimmune diseases (Lee *et al.*, 1994). Thus, inhibitor compounds capable of specifically inactivating certain critical kinases may have considerable therapeutic benefit in a number of clinical diseases.

All tyrosine and serine/threonine protein kinases have a region of approximately 300 amino acids known as the catalytic subunit which has evolved from a common ancestor kinase (Hanks and Quinn, 1991). Crystal structure determination of several kinases has shown that they all have a common bi-lobal structure (Wilson *et al.*, 1996; Zhang *et al.*, 1994; Xu *et al.*, 1997). The amino-terminal part of the subunit encodes a small lobe responsible for the binding of ATP, whereas the carboxy-terminal residues encode a larger lobe important for protein substrate binding. In the tertiary structure of the active kinase, both the ATP and the protein substrate binding sites are brought together allowing transfer of the ATP γ -phosphate to the amino acid acceptor on the protein substrate. The protein/peptide binding groove stretches across the face of the large lobe between two α -helices and under the small lobe. This groove therefore contains the residues important for defining the substrate specificity of the kinase.

Many protein kinases are arranged in kinase cascades within the cell, providing the ability for signal amplification in post-transduction pathways. This amplification relies on the upstream kinase specifically activating its downstream partner. For this reason, protein

kinases have developed remarkable substrate specificities which prevent unwanted crosstalk between different kinase cascades. This substrate specificity can be exploited in the design of selective protein kinase inhibitors.

A technique has recently been developed by L. Cantley's laboratory to provide, consensus peptide protein kinase substrate information (WO 95/18823, Songyang *et al*, 1994). First, a degenerate library of peptides with a phospho-acceptor such as tyrosine or serine/threonine flanked by amino acids on each side is synthesised. A preferred number of degenerate residues on each side of the phosphorylation site is four (corresponding to positions -1, -2, -3, -4, +1, +2, +3, +4) relative to the phosphorylated residue. Thus the library consists of peptides having a length of nine amino acids. The library is then phosphorylated by the protein kinase of interest and phosphorylated peptides isolated from the non-phosphorylated peptides by DEAE-sephacel and ferric chelation chromatography. The phosphopeptide mixture is then sequenced and the frequency of each amino acid at every position assessed to give a preferred substrate sequence. These studies have yielded consensus substrate information, but do not allow a detailed analysis of particular preferences for neighbouring residue interactions as pools of peptides are examined. Furthermore, this type of analysis may not show up rare good substrates which could be hidden by the presence of numerous poor substrates in the peptide pool. By this method individual peptides can never be identified as individual sequences, the result is that an average picture of substrate specificity is reached. Part of the problem is that each individual peptide is represented at such a low level, and many inevitably will not even be present. The results from Cantley's method do not represent individual peptides but a consensus picture of protein kinase substrate specificity.

Filamentous phage expressing gene III-linked degenerate peptide sequences have also been used to generate substrate information (Schmitz *et al*, 1996; Dente *et al*, 1997), however

this method is labour intensive and does not allow the use of unnatural amino acids or peptidomimetics. Substrate information can also be obtained from knowledge of the physiological kinase substrates. This approach is limited and previous attempts to utilise this information for the design of successful therapeutic cell permeable protein kinase inhibitors has failed (Kemp *et al.*, 1991).

We believe that identification of detailed substrate characteristics can intimately map the substrate binding groove and provide information that can lead to the design of enzyme inhibitor molecules. For the reasons described above, there are no current methods for obtaining this information. Therefore, we have invented a method of using small molecules, in a self deconvoluting library format, to probe a larger active site by positional scanning of a target group. This method is rapid, not labour intensive and results in the identification of discrete sequences.

Summary of Invention

This invention provides for the active site mapping of enzymes which catalyse covalent modification including, but not limited to phosphorylation, acylation, dephosphorylation in which a fixed residue (hereafter known as the catalytic residue) such as a tyrosine, serine, threonine, histidine, aspartic acid residue or any other residue containing an appropriate side chain is modified. The method of the invention has an additional level of complexity over and above that of the self-deconvoluting libraries described in WO97/42216 (the content of which is incorporated herein by reference, where legally permissible).

This involves making a library of smaller libraries (referred to as sub-sets) where a fixed residue is moved stepwise through the sequence of amino acids or other groups (such as

peptidomimetics [any compound that can be added to the substrate or inhibitor chain]]. The result is that in each sub-set of the library the fixed residue is found in a different position. For example, in a library using four variable positions, five sub-sets in each library have to be made, ZXXXX, XZXXX, XXZXX, XXXZX and XXXXZ where Z is the fixed residue and X are the four variable residues. We recognise that there may be a need in certain circumstances for further invariant residues, however these would occupy fixed positions and would not affect either the scanning or the self-deconvoluting of the libraries. The invariant residue(s) might be fixed in position relative to the modifiable residue Z or may be fixed in position relative to the overall motif sequence. Additional fixed residues can be added if desired, or one of the variable residues can be made invariant. In the later case the library would be a small part of the libraries described here. Cases where it is desirable to include one or more fixed residues include libraries required to look at enzymes which always require another invariant residue in another position. However, in cases where two fixed residues are required, and they are both modified, it can be desirable to include this residue in one of the variable positions (i.e. make it one of the residues chosen in a variable position). The reason for this is that the sequence of events (the order in which the two residues become modified) can then also be probed by this scanning library technique. In this case it may also be beneficial to make an additional library in which the fixed residue is not present at all, corresponding to the library XXXXX. We would therefore have a library of six sub-sets. These modifications are within the scope of the invention and would be recognised by someone skilled in the art.

It can be readily seen that by combining the data from each library sub-set, the residues from -4 to +4 either side of the catalytic residue can be mapped:

A-B-C-D-Z

B-C-D-Z-E

C-D-Z-E-F

D-Z-E-F-G

Z-E-F-G-H

The mapped sequence would therefore be A-B-C-D-Z-E-F-G-H.

The above example using 5 subsets of libraries of peptides of 5 amino acids allows the mapping of a sequence of 9 amino acids. In general one could carry out the invention using n subsets of n -mer peptides so as to provide mapping data for the residues from $-(n-1)$ to $+(n-1)$ either side of the active site. Thus in general the length of the mapped sequence would be $(2n)-1$.

Where the residue type at any given position relative to the fixed residue is similar in different subsets, the data can be used in an additive manner. For example, if an aromatic residue is required adjacent to the fixed residue, then any sequences which contain this feature in any of the library subsets can be considered in an additive way.

In this invention there is no need to separate modified from unmodified sequences because of the self deconvoluting nature of the library. The assay screen produces a series of hits, the patterns of which reveal the unique sequences in each well. This enables a pattern of substrate preferences to be determined for any enzyme.

The unique sequences obtained using this invention can be used to provide substrates for high throughput assays and provide detailed information about the active site to aid rational drug design.

This invention can also be used as an inhibitor library to screen against known modifying enzymes where a known substrate exists and can be set up in an assay format. Those skilled in the art would realise that by replacing the fixed residue with a suitable compound that is not modified an inhibitor library can be constructed. For example if a modifiable fixed tyrosine were to be changed to a tyrosine derivative residue that cannot be phosphorylated, such as halogenated tyrosine, dopamine, or tyrosine substituted by aromatic compounds, then an inhibitor library will be formed. This could allow the more direct identification of prototype inhibitors of enzymes for rational drug design.

Use of these libraries could be extended to other systems where a defined endpoint is desired, but the target enzyme is unknown. Such examples could include, but are not limited to, bacterial lysis in growing cultures or inhibiting phosphorylation of transcription factors in cell lysates.

In one embodiment of this invention the sequences identified by this method are considerably smaller than have previously been reported for library screens on protein kinase substrates, which makes them more amenable to computer modelling and drug design. Furthermore, this methodology provides information about the relative relationships between neighbouring residues of active substrates; information which is not available from a straightforward oriented degenerate peptide library approach used by Cantley (Songyang *et al*, 1994). Thus, this novel methodology provides a significant improvement in the quality of substrate based information that is achievable, in comparison to that produced from previously described methods.

This invention allows data to be obtained from single peptide rankings which could be used to rationally design sets of enzyme inhibitor molecules which compete with the physiological substrate for binding to the active site of the enzyme.

Description of the Drawings

Figure 1. Design of protein tyrosine kinase library. Each peptide consists of a biotin tag, an epsilon amino hexanoic acid spacer and 5 amino acids, including a phosphorylatable tyrosine residue. Each of the amino acid positions A-H is varied as described. For example, A1-10 means that position A is varied using 10 defined amino acids.

Figure 2. Best substrates identified by screening tyrosine library sub-sets 1 to 5 against ZAP-70 protein tyrosine kinase. The protein tyrosine kinase library described in Figure 1 was phosphorylated for 30 minutes at 30°C using the catalytic domain of human ZAP-70. Peptides were captured using streptavidin-coated 96 well plates and phosphotyrosine detected using anti-phosphotyrosine antibody, anti mouse IgG-HRP and tetramethylbenzidine (see experimental methods). Best substrates were identified as those which gave the highest amount of phosphate incorporation.

Figure 3. Km Determination of Biotin-εAHA-DEEDYFE(Nle) [SEQ ID NO. 3]. The catalytic domain of human ZAP-70 was used to phosphorylate varying concentrations of peptide for 10 minutes at 30°C in the presence of ^{33}P -γ-ATP. Peptide capture was performed using streptavidin filter plates, scintillation fluid added, and counting performed using a beta-counter (see experimental methods). Samples were assayed in triplicate.


Description of the Invention

This invention provides for the active site mapping of enzymes which catalyse covalent modification including, but not limited to, phosphorylation, acylation, dephosphorylation in which a residue such as a tyrosine, serine, threonine, histidine, aspartic acid or any other residue containing an appropriate side chain is modified.

Thus, the invention provides a method for determining an amino acid sequence motif or a peptidomimetic sequence motif containing an active site capable of being bound by an enzyme which catalyses covalent modification of a substrate molecule, comprising;

- a) contacting the enzyme with a library consisting of a number of oriented degenerate library subsets of molecules, each subset comprising unmodified degenerate motif sequences each having n residues and each having a modifiable residue at a different fixed non-degenerate position, under conditions which allow for modification of molecules which are a substrate for the enzyme;
- b) allowing the enzyme to modify modifiable residues in library subsets containing molecules having an active substrate site for the enzyme;
- c) deconvoluting the oriented degenerate library subsets of the library, *in situ* without separating modified from unmodified molecules, so as to reveal the sequence of any motif which has been modified by covalent binding of the enzyme;

wherein each library subset is of formula (I)



wherein

Zaa is a non-degenerate modifiable natural or unnatural amino acid residue or peptidomimetic;

Xaa is any natural or unnatural amino acid residue or peptidomimetic;

x and y are each independently 0 or an integer;

$(x + y) = (n-1)$; and

n = an integer from 3 to 8, preferably 5.

This invention can be applied for instance to a protein tyrosine kinase in order to exemplify the technology. It provides a rapid method of identifying discrete protein kinase substrate sequences which allows pharmacophore generation and design of active site inhibitors. This invention can also be used to directly identify protein kinase inhibitor molecules. General formula: (Xaa)_x Tyr (Xaa)_y [SEQ.ID No. 2].

In the first exemplification of this invention, a recombinant form of the human ZAP-70 enzyme was used in an *in vitro* phosphorylation reaction to phosphorylate the five substrate sub-libraries which scan the sequence -4 to +4 around a central tyrosine residue (Figure 1). The libraries were arranged in 96 well microtitre plate format with pools of 20 peptides in each microtitre well. However, those skilled in the art will realise that the library can be constructed on any scale. For example the library Sub-Set can be miniaturised on a "chip" scale or constructed on a large bulk scale depending on the requirements of the library.

Library peptides were made with biotin tags, which allowed peptide capture on streptavidin-coated microtitre plates. Detection of phosphotyrosine was achieved using anti-phosphotyrosine antibody detection in an ELISA assay using tetramethylbenzidine substrate and recording absorbance at 450 nm. Background absorbance readings of 0.1 to 0.2 were recorded while the highest substrate peptide value was 1.5. Deconvolution of the hit peptides was performed as described in WO 97/42216. Clear defined substrates were deconvoluted in library sub-sets 1 to 4, but not in 5. This probably reflects the absolute requirement of ZAP-70 for an amino acid residue in the -1 position.

For the purpose of this exemplification, the peptides used were tagged with d-Biotin and a linker (epsilon amino hexanoic acid or some other spacing group). In principle any tag and linker can be used, although this invention also provides that a tag and linker does not have to be present if mass spectroscopy, for example, is used to identify the peptide hits. The

purpose of the tag is solely to enable capture of all of the peptides (whether modified or not) so that excess reagents can be washed away. The reporting systems to detect peptide modification can include, but are not limited to, antibody recognition, radioactive assay or mass spectroscopy.

In the library used to exemplify the invention, a Biotin tag was chosen because we believed that this would give improved results. The reasoning for this choice of tag was because of the high level of positively charged groups on the enzyme in the area in which the tag sits. This charged area would cause unfavourable interactions with tags more commonly used by others in the field, such as poly-lysine or poly-arginine. We would expect this reasoning to be applicable to any enzyme which binds a highly negatively charged molecule such as but not limited to ATP, close to the peptide binding site.

Tags are preferably non peptidic, with as little charge, either positive or negative, as possible. Biotin is a good example of this. The aim is to minimise the interactions of the tag with the protein so the resultant hits are largely due to the binding of the peptides rather than reflecting the binding of the tag. The best method of all if this argument is applied to its logical conclusion would be to not use a tag at all and use mass spectroscopy to identify the peptides. However, currently this approach is of limited value due to the time taken to run and analyse a library of the size used here to exemplify the invention.

The results obtained from the library screen clearly demonstrated amino acid residues preferred by the protein kinase at each of the -4 to +4 sites (Figure 2). The 5 mer peptides overlapped to give information on amino acid preference at each of the binding positions -4 to +4. To confirm this a consensus peptide, Biotin-eAHA-DEEDYFE(Nle) [SEQ ID No. 3], representing the best -4 to +4 amino acids was made and tested as a substrate (Figure 3). This substrate gave a K_m against ZAP-70 of 15.79 μM , which is better than the best ZAP-

Word...39200PCT.JO-30th October 1998

12

70 substrate described in the literature, a longer peptide of 14 amino acids with a tag of 3 arginines and a K_m of 29 μ M (Wandenburg *et al*, 1996).

In the second application of this invention, a recombinant form of the human Syk enzyme was used in an *in vitro* phosphorylation reaction to phosphorylate the five substrate sub-libraries which scan the sequence -4 to +4 around a central tyrosine residue, as previously performed for the ZAP-70 library. Detection of phosphotyrosine was achieved using anti-phosphotyrosine antibody detection in an ELISA assay using tetramethylbenzidine substrate and recording absorbance at 450 nm. Background absorbance readings of 0.10 were recorded while the highest substrate peptide value was 1.46. Deconvolution of the hit peptides was performed as described in WO 97/42216. Clear defined substrates were deconvoluted in all library sub-sets.

In the third application of this invention, a recombinant form of the human CSK enzyme was used in an *in vitro* phosphorylation reaction to phosphorylate the five substrate sub-libraries which scan the sequence -4 to +4 around a central tyrosine residue, as previously performed for the ZAP-70 library. Detection of phosphotyrosine was achieved using anti-phosphotyrosine antibody detection in an ELISA assay using tetramethylbenzidine substrate and recording absorbance at 450 nm. Background absorbance readings of 0.04 were recorded while the highest substrate peptide value was 0.22. Deconvolution of the hit peptides was performed as described in WO 97/42216. Clear defined substrates were deconvoluted in all library sub-sets.

In the fourth application of this invention, a recombinant form of the Abelson murine leukaemia virus protein tyrosine kinase v-Abl was used in an *in vitro* phosphorylation reaction to phosphorylate the library sub-set 4 which scans the sequence -1 to +3 around a zero position tyrosine residue, as previously performed for the ZAP-70 library. Detection of

phosphorytyrosine was achieved using anti-phosphotyrosine antibody detection in an ELISA assay using tetramethylbenzidine substrate and recording absorbance at 450 nm. Background absorbance readings of 0.11 were recorded while the highest substrate peptide value was 0.32. Deconvolution of the hit peptides was performed as described in WO 97/42216. Clear defined substrates were deconvoluted in the library sub-set.

In the fifth application of this invention, the invention was used to map the substrate specificity of a protein serine or serine/threonine kinase (which include I-kappa B kinase beta and cAMP-dependent protein kinase [cAPK]). A protein serine or serine/threonine kinase enzyme was used in an *in vitro* phosphorylation reaction to phosphorylate the five substrate sub-libraries which scan the sequence -4 to +4 around a central serine residue. The library was synthesised as the protein tyrosine kinase ZAP-70 library save that the tyrosine fixed residues were replaced with a serine which was then scanned through the five sub-libraries. Detection of phosphoserine was achieved using anti-phosphoserine antibody detection in an ELISA assay using tetramethylbenzidine substrate and recording absorbance at 450 nm. Deconvolution of the hit peptides was performed as described in WO 97/42216.

General formula: (Xaa)_x Ser (Xaa)_y [SEQ ID No. 4].

Library Sub-Set 1 Xaa-Xaa-Xaa-Xaa-Ser

Library Sub-Set 2 Xaa-Xaa-Xaa-Ser-Xaa

Library Sub-Set 3 Xaa-Xaa-Ser-Xaa-Xaa

Library Sub-Set 4 Xaa-Ser-Xaa-Xaa-Xaa

Library Sub-Set 5 Ser-Xaa-Xaa-Xaa-Xaa

Likewise the Library Sub-Sets can be synthesised for the mapping of threonine kinases by the synthesis of a library containing the threonine residue to allow phosphorylation by enzymes recognising this residue.

It will be realised by those skilled in the art that the replacement of the recognition residue such as the tyrosine, or serine, with a residue that is covalently modified by the enzyme to be mapped allows the active site of any such enzyme to be determined according to the invention.

The invention will now be described by reference to the following examples.

Example 1

In this example the invention was used to map the active catalytic site of ZAP-70, a protein kinase enzyme that catalyses the phosphorylation of a tyrosine residue. The example illustrates the synthesis of a number of compounds, and their use as a sub-set library for the mapping of the enzyme so as to allow the subsequent identification and synthesis of single specific substrates for the enzyme.

Synthesis of Peptide Compounds.

Preparation of Crown Assembly

The peptide compounds were synthesised in parallel fashion using Fmoc-Rink-DA/MDA derivatised gears (ex Chiron Mimotopes, Australia) loaded at approximately 1.6 μ M per crown. Prior to synthesis each crown was connected to its respective stem and slotted into the 8 x 12 stem holder. Coupling of the amino acids employed standard Fmoc amino acid chemistry as described in 'Solid Phase Peptide Synthesis', E. Atherton and R.C. Sheppard, IRL Press Ltd, Oxford, UK, 1989.

Removal of N-Fmoc Protection

A 250 ml solvent resistant bath is charged with 200 ml of a 20% piperidine/DMF solution. The multipin assembly is added and deprotection allowed to proceed for 30 minutes. The assembly was removed and excess solvent removed by brief shaking. The assembly is then washed consecutively with (200 ml each), DMF (5 minutes) and MeOH (5 minutes, 5 minutes, 5 minutes) and left to air dry for 15 minutes.

Quantitative UV Measurement of Fmoc Chromophore Release

A 1 cm path length UV cell is charged with 1.2 ml of a 20% piperidine/DMF solution and used to zero the absorbance of the UV spectrometer at a wavelength of 290nm. A UV standard is then prepared consisting of 5.0 mg Fmoc-Asp(OBut)-Pepsyn KA (0.08 mmol/g) in 3.2 ml of a 20% piperidine/DMF solution. This standard gives $Abs_{290} = 0.55-0.65$ (at room temperature). An aliquot of the multipin deprotection solution is then diluted as appropriate to give a theoretical $Abs_{290} = 0.6$, and this value compared with the actual experimentally measured absorbance showing the efficiency of previous coupling reaction.

Coupling Of Standard Amino Acid Residues

Coupling reactions were performed by charging the appropriate wells of a polypropylene 96 well plate with the pattern of activated solutions required during a particular round of coupling. Gear (approx 1.6 μ mole) standard couplings were performed in DMF (300 μ l).

Coupling of an Amino-acid Residue To Appropriate Well

Whilst the multipin assembly is drying, the appropriate N-Fmoc amino acid pfp esters (10 equivalents calculated from the loading of each crown) and HOBt (10 equivalents) required for the particular round of coupling are accurately weighed into suitable containers. Alternatively, the appropriate N-Fmoc amino acids (10 equivalents calculated from the loading of each crown), desired coupling agent e.g. HBTU (9.9 equivalents calculated from the loading of each crown) and activation e.g. HOBt (9.9 equivalents calculated from the loading of each crown), NMM (19.9 equivalents calculated from the loading of each crown) were accurately weighed into suitable containers.

The protected and activated Fmoc amino acid derivatives were then dissolved in DMF (300 l for each gear e.g. for 20 gears, 20 x 10 eq. x 1.6 μ moles of derivative would be dissolved in 10 ml DMF). The appropriate derivatives were then dispensed to the appropriate wells ready for commencement of the 'coupling cycle'. As a standard, coupling reactions were allowed to proceed for 6 hours. The coupled assembly was then washed as detailed below.

Coupling of d-Biotin acid to pins

d-Biotin (10eq), 1-hydroxybenzotriazole. H_2O (10eq), BOP (9.95eq) and NMM (19.9eq) were dissolved in DMF (0.3mL per well) and agitated for 2 minutes. 300 μ L of solution was dispensed to each well of a 96-well polypropylene plate. The gears were then added to the solution and left for 24 hours. Fresh solution was made up, the gears washed in DMF for 5 minutes and then added to the fresh coupling mixture and left a further 24 hours.

The pin assembly was removed from the plate, shaken free of excess liquid then immersed in DMF (200mL) for 5mins. The assembly was again shaken then immersed in MeOH (200mL, 3 x 5mins) and allowed to air dry.

Washing Following Coupling

If a 20% piperidine/DMF deprotection is to immediately follow the coupling cycle, then the multipin assembly is briefly shaken to remove excess solvent washed consecutively with (200 ml each), MeOH (5 minutes) and DMF (5 minutes) and de-protected (see 6.2). If the multipin assembly is to be stored or reacted further, then a full washing cycle consisting brief shaking then consecutive washes with (200 ml each), DMF (5 minutes) and MeOH (5 minutes, 5 minutes, 5 minutes) is performed.

Following these general methods, the peptide libraries shown in Figure 1 were sequentially assembled by applying the appropriate coupling procedure at the correct cycle during synthesis.

Acidolytic Mediated Cleavage of Peptide-Pin Assembly

Acid mediated cleavage protocols were strictly performed in a fume hood. A polystyrene 96 well plate (1 ml/well) was labelled, then the tare weight measured to the nearest mg. Appropriate wells were charged with a trifluoroacetic acid/triisopropylsilane (95:5, v/v, 600 μ l) cleavage solution, in a pattern corresponding to that of the multipin assembly to be cleaved.

The multipin assembly is added, the entire construct covered in tin foil and left for 2 hours. The multipin assembly is then added to another polystyrene 96 well plate (1 ml/well) containing trifluoroacetic acid/triethylsilane (95:5, v/v, 600 μ l) (as above) for 5 minutes.

Work up of Cleaved Peptides

The primary polystyrene cleavage plate (2 hour cleavage) and the secondary polystyrene plate (5 minute wash) were then placed in the SpeedVac and the solvents removed (minimum drying rate) for 90 minutes.

The contents of the secondary polystyrene plate were transferred to their corresponding wells on the primary plate using an acetonitrile/water/acetic acid (50:45:5, v/v/v) solution (3 x 150 µl) and the spent secondary plate discarded.

Analysis of Products

A 5µL aliquot from each well is diluted to 100 µl with 0.1% aq. TFA, then a 10µL aliquot from this plate diluted with a further 100 µl 0.1% aq. TFA. The double diluted plate was analysed by HPLC-MS.

Final Lyophilisation of Peptides

The plate was covered with tin foil, held to the plate with an elastic band. A pin prick was placed in the foil directly above each well and the plate placed at -80°C for 30 minutes. The plate was then lyophilised on the 'Heto freeze drier' overnight. Finally, the dried plate was weighed. The total cleaved peptide was quantified (by weight) and the average content of each peptide calculated. Since all the peptides present have originated from the same peptide-pin assembly, cleaved under identical conditions, it is reasonable to assume that the contents of each well are roughly equimolar.

Protein kinase cloning, expression and purification

Polymerase chain reaction (PCR) and downstream cloning

The coding sequence for human ZAP-70 amino acid 306-615 was amplified from Jurkat T cell cDNA by PCR (2 minutes at 94°C, followed by 35 cycles of 15 seconds 94°C, 30 seconds 65°C, 2 minutes 72°C and a final single 5 minute 72°C incubation) using the primers:

5' CCGGGATCCGCCATGCCCATGGACACGAGCGTGTAT 3' [SEQ ID No. 5]

5' GGGGGATCCTCAGTGGTGGTGGTGGTGGTGGGCACAGGCAGCCTCAGC
CTTCTGTGT 3' [SEQ ID No. 6]

The PCR amplicon was cloned into the Bam H1 site of pUC19 and sequence confirmed using M13-20 and reverse primers on an Applied Biosystems Prism 310 sequencer as described by manufacturer. The Bam H1 ZAP-70 insert was excised from the sequencing vector and ligated into the Baculoviral transfer vector pAcUW51 (Pharmingen).

Generation of ZAP-70 enzyme using baculovirus

Homologous recombination with wild type baculoviral DNA was then performed in Sf9 insect cells and viral supernatant harvested. Plaque purified virus was exposed to several viral amplification steps then used at a titre of 3×10^9 PFU/ml to infect 3×10^6 cells/ml Sf9 cells at an MOI of 10 in an Applecon bioreactor using 60% dissolved oxygen. Cells were harvested 3 days post infection.

Protein purification

The infected cell pellet was lysed in 50 mM Tris-HCl, pH 7.5, 0.15 M NaCl, 25% sucrose, 1 mM 4-nitrophenol phosphate, 1 mM sodium orthovanadate and protease inhibitors.

Following homogenisation using a dounce pestle B, the cleared lysate was loaded onto a cobalt-sepharose column. After column washing with lysis buffer, elution was performed with an imidazole gradient and ZAP-70 fractions identified by protein kinase activity against the peptide substrate Lys-Lys-Lys-Lys-Ala-Asp-Glu-Glu-Asp-Tyr-Phe-Ile-Pro-Pro-Ala as described in Casnellie *et al*, 1991

Library Screening

Library peptides were phosphorylated in pools of 20 peptides at a final concentration of 1 μ M total peptide in 50 mM HEPES, pH 7.5, 0.1% Triton X-100, 100 μ M ATP, 10 mM $MnCl_2$, 1 mM DTT and 0.2 mM sodium orthovanadate for 30 minutes at 30°C. These reaction mixtures were then stopped using 100 mM EDTA, 6 mM adenosine, transferred to streptavidin-coated microtitre plates and allowed to bind for 30 minutes at 20°C. Unincorporated reaction products were washed from the plate using PBS/0.1% Tween 20 then plate incubation performed with anti-phosphotyrosine antibody (Sigma mouse monoclonal clone PT66) in 2% BSA/PBS/Tween for 1 hour at 20°C. Unbound antibody was removed by plate washing using PBS/Tween then incubation performed with rabbit anti mouse-HRP (Amersham) for a further 1 hour. HRP detection was then performed with tetramethylbenzidine (TMB) substrate and measuring absorbance at 450 nM using a spectrophotometer (Dynex MRX).

The best substrates were identified as those which gave the highest amount of phosphate incorporation. The library subsets were deconvoluted according to the teaching of WO97/42216: this gives an immediate determination of the unique sequence of any phosphorylated motif without the need for further synthesis or sequencing. (Figure 2 [SEQ ID Nos. 7,8,9,10]).

Peptide K_m determination

Biotin-tagged peptides were phosphorylated at varying concentrations in 50 mM HEPES, pH 7.5, 0.1% Triton X-100, 200 μ M ATP, 10 mM $MnCl_2$, 1 mM DTT and 0.2 mM sodium orthovanadate using 0.5 μ Ci ^{33}P - γ -ATP for 10 minutes at 30°C. The reactions were stopped using 2 M guanidine hydrochloride, diluted 1 in 10 in water then 5 μ l reaction mix spotted onto SAMTM titre plates (Promega). Unincorporated reaction products were washed as described by manufacturer then 20 μ l scintillation liquid added and plate counted on a Packard TopCount beta-counter. K_m was calculated using a non-linear one site hyperbola model (ATP was added in excess to negate influence of the ATP binding site on the substrate site kinetics). (Figure 3).

Example 2

In this example the invention was used to map the active catalytic site of Syk, a protein kinase enzyme that catalyses the phosphorylation of a tyrosine residue. The example illustrates the positional scanning of the sub-set libraries for the mapping of the enzyme, and their use so as to allow the subsequent identification of the preferred substrates of the enzyme catalytic site.

The mapping and assessment of the catalytic site was performed as detailed in Example 1. The substrate preferences were deconvoluted as detailed in WO 97/42216 and are detailed below.

Library Sub-Set 1	Asp-Glu-Glu-Asp-Tyr [SEQ ID No. 11]
Library Sub-Set 2	Asp-Glu-Glu-Tyr-Asp [SEQ ID No. 12]

22

Library Sub-Set 3	Asp-Glu-Tyr-Glu-Asp [SEQ ID No. 13]
Library Sub-Set 4	Asp-Tyr-Glu-Glu-Val [SEQ ID No. 14]
Library Sub-Set 5	Tyr-Ser-Ile-Ile-Nle [SEQ ID No. 15]

Example 3

In this example the invention was used to map the active catalytic site of CSK. The subset library was used to scan the enzyme active site so as to allow the subsequent identification and synthesis of the preferred specific substrates for the enzyme as listed below.

Library Sub-Set 1	Asp-Glu-Glu-Glu-Tyr [SEQ ID No. 16]
Library Sub-Set 2	Asp-Glu-Glu-Tyr-Phe [SEQ ID No. 17]
Library Sub-Set 3	Asp-Glu-Tyr-His-Asn [SEQ ID No. 18]
Library Sub-Set 4	Asp-Tyr-His-Leu-Phe [SEQ ID No. 19]
Library Sub-Set 5	Tyr-Pro-Ile-Glu-Val [SEQ ID No. 20]

Example 4

In this example the invention was used to map the active catalytic site of v-Abl, a protein kinase enzyme that catalyses the phosphorylation of a tyrosine residue. For this enzyme only the Library Sub-Set 4 (i.e. X-Tyr-X-X-X according to SEQ ID No. 2) was scanned with the enzyme. The active site substrate recognition substrate for this enzyme for this Sub-Set was Serine-tyrosine-phenylalanine-histamine-glutamine [SEQ ID No. 21].

References

Arpaia E, Shahar M, Dadi H, Cohen A and Roifman CM. (1994). Defective T cell receptor signaling and CD8+ thymic selection in humans lacking zap-70 kinase. *Cell* 76(5):947-958

Barker KT, Jackson LE, and Crompton MR. (1997). BRK tyrosine kinase expression in a high proportion of human breast carcinomas. *Oncogene* 14;15(7):799-805

Casnellie, JE. (1991). Assay of protein kinases using peptides with basic residues for phosphocellulose binding. *Methods Enz.* 200, 115-120.

Denti, L., Vetriani, C., Zucconi, A., Pelicci, G., Lanfranccone, L., Pelicci, P. G. and Cesareni, G. (1997). Modified phage peptide libraries as a tool to study specificity of phosphorylation and recognition of tyrosine containing peptides. *J. Mol. Biol.*, 269, 694-703.

Elder ME, Lin D, Clever J, Chan AC, Hope TJ, Weiss A, Parslow TG. (1994). Human severe combined immunodeficiency due to a defect in ZAP-70, a T cell tyrosine kinase. *Science* 264(5165):1596-1599

Hanks, SK and Quinn, M. (1991). Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members. *Methods Enzymol* 200, 38-62

Kemp, BE, Pearson, RB, and House, CM, (1991). Pseudosubstrate-based peptide inhibitors. *Methods Enzymol* 201, 287-304.

Lee JC, Laydon JT, McDonnell PC, Gallagher TF, Kumar S, Green D, McNulty D, Blumenthal MJ, Heys JR, Landvatter SW, *et al.* (1994). A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature* 372(6508):739-746

Lehtola L, Partanen J, Sistonen L, Korhonen J, Warri A, Harkonen P, Clarke R and Alitalo K. (1992). Analysis of tyrosine kinase mRNAs including four FGF receptor mRNAs expressed in MCF-7 breast-cancer cells. *Int J Cancer* 50(4):598-603

Roifman CM. (1995). A mutation in zap-70 protein tyrosine kinase results in a selective immunodeficiency. *J Clin Immunol* 15(6 Suppl):52S-62S

Schmitz, R., Baumann, G. and Gram, H. (1996). Catalytic specificity of phosphotyrosine kinases Blk, Lyn, c-Src and Syk as assessed by phage display. *J. Mol. Biol.*, 260, 664-677.

Songyang Z, Blechner S, Hoagland N, Hoekstra MF, Piwnicka-Worms H and Cantley LC. (1994). Use of an oriented peptide library to determine the optimal substrates of protein kinases. *Curr Biol.* 4(11):973-982

Wardenburg, J.B., Fu, C., Jackman, J.K., Flotow, H., Wilkinson, S.E., Williams, D.H., Johnson, R., Kong, G., Chan, A.C. and Findell, P. R. (1996). Phosphorylation of SLP-76 by the ZAP-70 protein-tyrosine kinase is required for T-cell receptor function. *J. Biol. Chem.* 271(33), 19641-19644.

Wilson KP, Fitzgibbon MJ, Caron PR, Griffith JP, Chen W, McCaffrey PG, Chambers SP and Su MS. (1996). Crystal structure of p38 mitogen-activated protein kinase. *J Biol Chem.* 271(44):27696-27700

Xu W, Harrison SC and Eck MJ. (1997). Three-dimensional structure of the tyrosine kinase c-Src. *Nature* 385(6617):595-602

Zhang F, Strand A, Robbins D, Cobb MH and Goldsmith EJ. (1994) Atomic structure of the MAP kinase ERK2 at 2.3 Å resolution. *Nature* 367(6465):704-711

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000